


**RESTRICTED USE PESTICIDE**

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

**Section 18  
EMERGENCY  
EXEMPTION****FOR DISTRIBUTION AND  
USE ONLY IN  
NORTH CAROLINA****Insecticide/Miticide****For Agricultural or Commercial Use Only**  
EPA File Symbol: 06-NC-02**EMERGENCY CALLS: 800-331-3148****IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN  
A MANNER INCONSISTENT WITH ITS LABELING****THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE  
APPLICATION.****This Specific Exemption is Effective from March 27, 2006 thru October 31, 2006.**

<b>Crop</b>	<b>Pest</b>	<b>Rate of Application</b>
Sweet Potato	Beetle Complex to include the adult and immature life stages of Flea beetles, click beetles (wireworms), cucumber beetles (rootworm), whitefringed beetles, and May/June beetles (white grub)	See Directions for Use for specific information

**DIRECTIONS FOR USE:**

Capture may be applied as an in-furrow or incorporated planting time treatment for the control of wireworms, rootworms and white grubs. Apply Capture at the rate of 0.3 pounds active per acre (19.2 ounces formulated) as an incorporated bed spray, in-furrow spray or T-band spray at planting time. Capture can then be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Capture to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Capture as a banded spray over the row at a rate of 0.05 pounds active per acre (3.2 ounces formulated) in 10 gallons per acre of spray.

Alternatively, Capture may be applied at the rate of 0.2 pounds active per acre (12.8 ounces formulated) as an incorporated bed spray, in-furrow spray or T-band spray followed up by a lay-by treatment of 0.15 pounds active per acre (9.6 ounces formulated) applied as a banded spray to the drill area covered up with soil utilizing cultivation equipment set to throw soil to the drilled area for control of wireworms, rootworms, and white grubs in 10 gallons per acre spray.

Capture may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles and May/June beetles (white grubs). Capture should be applied at the rate of 0.05 lbs active per acre (3.2 ounces formulated) in 10 gallons of spray by ground. Do not make more than 2 foliar applications per season

**RESTRICTIONS:**

Do not apply more than 0.5 pounds active ingredient per season.  
Do not apply within 30 days of harvest.  
Treated fields may not be grazed.  
Observe all the restrictions of the Capture 2EC label (EPA Reg. No. 279-3069).

**CONDITIONS OF USE:**

FMC Corporation intends that this Section 18 Emergency Exemption label be distributed only to end users and/or growers and/or applicators acting on behalf of growers who agree that FMC would not be held responsible for failure to perform and/or any crop damage from use of Capture 2EC for beetle complex control on sweet potato. If such terms and conditions are unacceptable, FMC requests the return of Capture 2EC at once unopened. FMC considers the user to have accepted such terms and conditions upon use of this product.

**ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL (EPA Reg. No. 279-3069) ARE TO BE FOLLOWED.****PERSONAL PROTECTIVE EQUIPMENT AND REENTRY INTERVALS ON THE BASIC LABEL MUST BE FOLLOWED.****THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Capture2EC\_18\_NC\_sweet potato\_3-27-2006.doc

Capture and **FMC** -FMC Trademarks  
©2006 FMC Corporation