FIFRA Section 24(c) Special Local Need



FLUMIOXAZIN GROUP 14 HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA



EPA Reg. No. 59639-99 EPA SLN No. NC-150007

FOR CONTROL OF WINTER ANNUAL BROADLEAF WEEDS IN CLARY SAGE

This label is valid until December 31, 2030 or until otherwise amended, withdrawn, cancelled or suspended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered container product label for Valor® SX Herbicide.

This labeling must be in the possession of the user at the time of application.

FOR WEED CONTROL IN CLARY SAGE GROWN FOR THE ESSENCE AND ESSENTIAL OIL INDUSTRY (Do not use treated clary sage for food or feed)

PREPLANT APPLICATION FOR RESIDUAL CONTROL OF WINTER ANNUAL BROADLEAF WEEDS IN CLARY SAGE

- Apply 1.0 to 2.0 oz/A of *Valor* SX Herbicide as a preplant application when applied 30 days before planting clary sage with soils that contain less than 80% sand.
- If soils contain greater than 80% sand, apply 1 oz/A *Valor* SX Herbicide 30 days before planting clary sage or 2.0 oz/A 45 days before planting.
- When tank mixing, use 1.0 oz/A for enhanced burndown and limited residual weed control.
- Use the 2.0 oz/A rate for residual weed control through planting.
- Apply 3.0 oz/A of *Valor* SX Herbicide as a preplant application when applied 45 or more days before planting clary sage.
- Sequential preplant applications are prohibited.

Application Restrictions

- Treated soil that is splashed onto newly emerged sage may result in temporary crop injury.
- A minimum of 0.5 inch of rainfall/irrigation must pass between application and planting of clary sage.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- Apply in a minimum of 15 GPA of water to ensure uniform soil coverage and adequate coverage of emerged weeds.
- Do not apply more than 3.0 oz/A of *Valor* SX Herbicide preplant.
- Do not apply more than 7.0 oz of *Valor* SX Herbicide per acre per season.

POSTEMERGENCE APPLICATION FOR RESIDUAL CONTROL OF WINTER ANNUAL BROADLEAF WEEDS IN CLARY SAGE

 Apply 1.0 to 2.0 oz/A of Valor SX Herbicide as a postemergence application when clary sage is a minimum of 4 inches in diameter.

Application Restrictions

- Valor SX Herbicide can be applied postemergence to clary sage up to the time of flower bud initiation.
- Do not apply Valor SX Herbicide to clary sage after it begins flower bud initiation.
- Do not apply more than 2 oz of *Valor* SX Herbicide per acre during a single application.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- Do not tank mix *Valor* SX Herbicide with any adjuvant, fertilizer or pest control product or severe injury to clary sage will occur.
- Apply in a minimum of 15 GPA of water to ensure adequate coverage of emerged weeds.
- Do not apply more than 7.0 oz of *Valor* SX Herbicide per acre per season.

Please contact Valent U.S.A. LLC at 1-800-6-VALENT (682-5368) if you have questions regarding use of this product.

FIFRA 24(c) Registrant: Valent U.S.A. LLC

P.O. Box 5075

San Ramon, CA 94583

Valor is a registered trademark of Valent U.S.A. LLC.