

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

CYHALOFOP-BUTYL

GROUP

1

HERBICIDE

Clincher® SF

EPA Reg. No. 62719-357

SLN No. NC-260003

For Distribution and Use Only in the State of North Carolina

Postemergence Weed Control in Carolina Gold, Charleston Gold, and Santee Gold Rice

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Clincher® SF before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of Clincher SF according to this FIFRA 24(c) Special Local Need Label is subject to all use precautions and limitations imposed by the label affixed to the container for Clincher SF.

Directions for Use

Refer to Clincher® SF product label for General Use Precautions, Mixing, and Application instructions.

Apply Clincher SF herbicide at 13.5-15 fl oz/acre by air in a minimum of 10 gallons/per acre. Use of an agriculturally approved crop oil concentrate or methylated seed oil adjuvant at a rate of 1 quart/acre must be used for all applications of Clincher SF. Read and follow all use directions and precautions on adjuvant labels. See main Clincher SF label for information on drift reduction.

Target Weeds and Application Rate:

Target Weeds	Application Rate and Stage of Grass Weed Development	
	13.5 fl oz/acre	15 fl oz/acre
Weeds Controlled	preflood, up to 4 leaf	post flood, prior to grass weed heading
Amazon (tighthead)		
sprangletop		
barnyardgrass		
bearded sprangletop		
broadleaf sprangletop		
fall panicum		
johnsongrass (seedling)		
junglerice		
red sprangletop		
Weeds Suppressed		
goosegrass		
large crabgrass		

Use Restrictions

- If applied to heading grasses, heavy weed densities and/or previous untreated areas (salvage treatment), only partial control should be expected. Regrowth of these grasses may occur.
- Allow 10 days or more between applications.
- Do not apply more than 15 fl oz in a single application.
- Do not apply more than 25 fl oz per year in both the first and ratoon crops combined.
- Do not apply near desirable vegetation such as corn, sugar cane, sudangrass, sorghum, grass grown for seed, millet, sod farms, stone, pome, citrus, cereals, grass, and all other desirable crops.
- Do not apply within 60 days of rice harvest.
- Do not rotate treated land to crops other than rice for 3 months following application.

24(c) Registrant:

**Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054 USA**

TM® Trademarks of Corteva Agriscience and its affiliated companies

R161-006

Accepted: 7/07/2026

Initial printing.