

# Section 24(c) Special Local Need Label

FOMESAFEN GROUP 14 HERBICIDE

#### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA

Reflex® Herbicide
For Control of Yellow Nutsedge in Pine Seedling Nurseries Only

EPA Reg. No. 100-993 EPA SLN No. NC-950010

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Section 24(c) label and main EPA-registered Reflex Herbicide label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this 24(c) label and the main EPA-registered Reflex Herbicide (EPA Reg. No. 100-993) product label.
- Failure to follow the directions for use and precautions on this label may result in poor

**PESTS:** Yellow nutsedge (*Cyperus esculentus*)

This treatment should be used as part of an overall integrated pest management (IPM) program. A single application of Reflex Herbicide will not eliminate yellow nutsedge. This herbicide can be used as part of an integrated program to keep nutsedge weak and suppressed.

#### **RESTRICTIONS**

- Do not apply a preemergence application and a postemergence application to the same area during the same year.
- Do not exceed a total of 2 pints per acre per year.
- Do not apply this product through any type of irrigation system.

**PREEMERGENCE APPLICATION:** Apply Reflex Herbicide at 2 pints per acre. Make application immediately after seeding as a preemergence spray in 30 to 40 gallons of water per acre. Apply mulch for seed cover. For best results, irrigate with ½ inch of water immediately after application.

**POSTEMERGENCE APPLICATION:** Apply Reflex Herbicide at 1.5 pints per acre as a postemergence spray in 30 to 40 gallons of water per acre. Make application after pine

seedlings are 8 weeks old (i.e., 8 weeks after seedling emergence.) or later. **DO NOT** use a spray adjuvant (i.e., crop oil or surfactant, etc.) with Reflex Herbicide as this may cause severe pine seedling injury. Reflex Herbicide may cause temporary chlorotic (yellow) spots on pine seedlings. Scout field and apply to nutsedge patches. For best results, irrigate with ½ inch of water within one week of application

## ©2023 Syngenta

Reflex® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419-8300

Label Code: NC0993008AA0503