



## North Carolina



Tobacco Trust Fund Commission

## List of Plant-Parasitic Nematodes Recorded in North Carolina

Weimin Ye

North Carolina's agricultural industry, including food, fiber, ornamentals and forestry, contributes \$84 billion to the state's annual economy, accounts for more than 17 percent of the state's income, and employs 17 percent of the work force. North Carolina is one of the most diversified agricultural states in the nation. Approximately, 50,000 farmers grow over 80 different commodities in North Carolina utilizing 8.2 million of the state's 12.5 million hectares to furnish consumers a dependable and affordable supply of food and fiber. North Carolina produces more tobacco and sweet potatoes than any other state, ranks second in Christmas tree and third in tomato production. The state ranks ninth nationally in farm cash receipts of over \$10.8 billion (NCDA&CS Agricultural Statistics, 2017).

Plant-parasitic nematodes are recognized as one of the greatest threat to crops throughout the world. Nematodes alone or in combination with other soil microorganisms have been found to attack almost every part of the plant including roots, stems, leaves, fruits and seeds. Crop damage caused worldwide by plant nematodes has been estimated at \$US80 billion per year (Nicol et al., 2011). All crops are damaged by at least one species of nematode. Most plant-parasitic nematodes live in soil and damage plants by feeding in large numbers on roots, impairing the plant's ability to take up water and nutrients. Severe root damage caused by nematodes typically results in aboveground symptoms that may include stunting, yellowing of leaves, incipient wilt, loss of plant vigor and/or an overall general decline in plant performance. Damage is often more pronounced when plants are under stress from lack of water or nutrients or when damaged by other diseases or insects. Although nematodes rarely kill plants, they can drastically limit plant growth and yields. Plant-parasitic nematodes are usually confined to localized areas in soil and spread very slowly by their own power. However, they may be dispersed more rapidly by movement of infested soil through cultivation, on soil clinging to farming tools and tillers, in water, wind or on roots of transplants.

NCDA&CS  
AGRONOMIC DIVISION  
NEMATODE ASSAY SECTION

PHYSICAL ADDRESS  
4300 REEDY CREEK ROAD  
RALEIGH NC 27607-6465

MAILING ADDRESS  
1040 MAIL SERVICE CENTER  
RALEIGH NC 27699-1040

PHONE: 919-733-2655  
FAX: 919-733-2837

DR. WEIMIN YE  
NEMATOLOGIST

DR. COLLEEN HUADAK-WISE  
DIVISION DIRECTOR

STEVE TROXLER  
AGRICULTURE COMMISSIONER

In the state of North Carolina, nematodes are a major threat to most crops. About 94 plant-parasitic nematode species have been recorded. Among them, root knot, soybean cyst, reniform, sting, lesion, lance, stubby root, tobacco cyst, ring, and foliar nematodes are considered the most important. The list of plant-parasitic nematodes in Table 1 is based in part on the nematode section of the North Carolina Plant Disease Index by K. R. Barker (Grand, 1985) and includes all recent research findings (Ye et al., 2012, 2013, 2015a, b; Ye, 2017; Zeng et al., 2012b, 2015; Holguin et al., 2015a, 2016) and nematode assay services provided by the NCDA&CS. Nematode entries in this list do not necessarily indicate that all nematodes associated with a given plant species are pathogenic to that plant species. Although all of these nematodes are obligate parasites, some of them often cause little or no damage. Furthermore, nematodes are primarily soil inhabitants and may feed on associated weeds and grasses. In a few instances, the nematodes included under a given plant species probably were feeding primarily on these associated plants/weeds or had fed on previous crops.

**Table 1. List of plant-parasitic nematodes recorded in North Carolina.**

No.	Nematode species	Crop or plant	References
1	<i>Anguina tritici</i>	Rye, wheat	Grand, 1985
2	<i>Aorolaimus</i> spp.	Camellia	Grand, 1985
3	<i>Aphelenchoides fragariae</i>	Ornamentals, strawberry	McCuiston et al., 2007; Kohl et al., 2010
4	<i>A. myceliophagus</i>	Mushrooms, turfgrass	Zeng et al., 2012b
5	<i>A. parietinus</i>	Cotton	Steiner, 1938; Society of Nematologists, 1984
6	<i>A. ritzemabosi</i>	Florist's daisy	Strider, 1979; Grand, 1985
7	<i>A. subtenuis</i>	Unknown	Society of Nematologists, 1984
8	<i>Belonolaimus gracilis</i>	Corn, cotton	Ruehle and Sasser, 1962
9	<i>B. longicaudatus</i>	Bermuda grass, common bean, corn, cotton, creeping grasses, peanut, soybean, tomato	Zeng et al., 2012b
10	<i>B. maritimus</i>	American beachgrass	Grand, 1985
11	<i>Bursaphelenchus xylophilus</i>	Black pine, loblolly pine, longleaf pine, pond pine, shortleaf pine, slash pine, Virginia pine	Ye & Giblin-Davis, 2013
12	<i>Cactodera weissi</i>	Bermuda grass, knotweed, tobacco	Grand, 1985; Ye, 2012
13	<i>C. sphagni</i>	Trees	Ruehle, 1968
15	<i>Ditylenchus dipsaci</i>	Alfalfa, ornamentals, white clover, wild onion	Barker and Sasser, 1959; Grand, 1985
16	<i>D. triformis</i>	Iris	Hirschmann and Sasser, 1955
17	<i>Dolichodorus heterocephalus</i>	Bermuda grass, camellia	Zeng et al., 2012b
18	<i>D. marylandicus</i>	Pine	Lewis and Golden, 1981
20	<i>Globodera tabacum solanacearum</i>	Carolina horsetail, tobacco	Melton et al., 1991
21	<i>Helicotylenchus dihystera</i>	Bermuda grass, common bean, corn, cotton, creeping grasses, loblolly pine, peach, peanut, slash pine, soybean	Aycock et al., 1976; Schmitt and Barker, 1988; Zeng et al., 2012b
22	<i>H. erythraeae</i>	Grass	Ruehle & Sasser, 1962
23	<i>H. nannus</i>	Loblolly pine, slash pine	Ruehle and Sasser, 1962
24	<i>Hemicaloosia graminis</i>	Turfgrass	Zeng et al., 2012a

<b>25</b>	<i>Hemicriconemoides chitwoodi</i>	Camellia, turfgrass	Ye and Robbins, 2000; López et al., 2012a; Zeng et al., 2012b
<b>26</b>	<i>H. wessoni</i>	Bermuda grass	Ye and Robbins, 2000; Zeng et al., 2012b
<b>27</b>	<i>Hemicycliophora conida</i>	Creeping bentgrass	Zeng et al., 2012b
<b>28</b>	<i>H. gigas</i>	Forest soil	Thorne, 1955
<b>29</b>	<i>H. gracilis</i>	Grass	Thorne, 1955
<b>30</b>	<i>H. mettleri</i>	Trees	Brzeski, 1974
<b>31</b>	<i>H. parvana</i>	Turfgrass	López et al., 2013; van den Berg et al., 2018
<b>32</b>	<i>H. robbinsi</i>	Turfgrass	López et al., 2013; Subbotin et al., 2014; van den Berg et al., 2018
<b>33</b>	<i>H. sheri</i>	Grass	Society of Nematologists, 1984
<b>35</b>	<i>Heterodera cyperi</i>	Ornamentals	Golden, et al., 1962
<b>36</b>	<i>H. glycines</i>	Common bean, soybean	Schmitt and Barker, 1988; Koenning, 2004
<b>37</b>	<i>H. lespedezae</i>	Bush clover	Golden & Cobb, 1963
<b>38</b>	<i>H. trifolii</i>	Daylilly, white clover	Grand, 1985; Ye, 2012
<b>39</b>	<i>Hoplolaimus columbus</i>	Cotton, soybean	Sher, 1963; Schmitt and Barker, 1988; Holguin et al., 2015a, b, 2016
<b>40</b>	<i>H. coronatus</i>	Loblolly pine, slash pine	Ruehle and Sasser, 1962
<b>41</b>	<i>H. galeatus</i>	Bermuda grass, boxwood, Chinese holly, corn, cotton, creeping bentgrass, creeping grasses, slash pine, soybean, tall fescue, white clover	Schmitt and Barker, 1988; Zeng et al., 2012b; Holguin et al., 2015a
<b>42</b>	<i>H. stephanus</i>	Soybean	Sher, 1963; Holguin et al., 2015a
<b>43</b>	<i>Longidorus brevianulatus</i>	Turfgrass	Society of Nematologists, 1984
<b>48</b>	<i>Meloidodera floridensis</i>	Azalea, grass, loblolly pine, slash pine	Ruehle and Sasser, 1962; Triantaphyllou and Hirschmann, 1973; Ye, 2012
<b>49</b>	<i>Meloidogyne arenaria</i>	Asparagus, azalea, carrot, daylily, peanut, corn, sweet potato, tobacco, soybean, tomato	Winstead & Sasser, 1956
<b>50</b>	<i>M. carolinensis</i>	Blueberry	Eisenback, 1982
<b>51</b>	<i>M. enterolobii</i>	Cotton, horseweed, morning glory, sicklepod, soybean, sweet potato, tobacco	Ye et al., 2013
<b>52</b>	<i>M. graminis</i>	Bermuda grass, blue oat grass, centipedegrass, creeping bentgrass, creeping grasses, meadow fescue, St. Augustine grass	Ye et al., 2015b; Zeng et al., 2012b
<b>53</b>	<i>M. hapla</i>	Boxwood, cabbage, cantaloupe, common bean, cotton, gingseng, Irish potato, pea, peanut, peony, sage, soybean, strawberry, sweet potato, tobacco, tomato, watermelon, wheat	Schmitt and Barker, 1988; Society of Nematologists, 1984

<b>54</b>	<i>M. incognita</i>	Alfalfa, asparagus, azalea, bean, begonia, bentgrass, bermuda grass, boxwood, Buddleja, butterbean, cabbage, camellia, cantaloupe, carrot, collards, common bean, corn, cotton, creeping bentgrass, creeping grasses, cucumber, daylily, dogwood, eggplant, English oak, fig, forsythia, gardenia, green bean, holly, hyacinth bean, hydrangea, kiwifruit, Korean boxwood, lantana, liriope, lima bean, mondo grass, milo, muskmelon, oak, okra, ornamentals, peach, pepper, peony, potato, pumpkin, sage, snap bean, soybean, spinach, squash, St. Augustine grass, strawberry, sweet potato, tobacco, tomato, tube rose, watermelon, white clover, zoysiagrass	Schmitt and Barker, 1988
<b>55</b>	<i>M. javanica</i>	Boxwood, southern peas, soybean, sweet potato, tobacco, tomato	Schmitt and Barker, 1988
<b>56</b>	<i>M. marylandi</i>	Bermuda grass, creeping bentgrass, creeping grasses	Ye et al., 2015b
<b>57</b>	<i>M. megatyla</i>	Loblolly pine	Baldwin and Sasser, 1979
<b>58</b>	<i>M. naasi</i>	Bermuda grass, creeping bentgrass, creeping grasses	Zeng et al., 2012b; Ye et al., 2015b
<b>60</b>	<i>M. spatinae</i>	Smooth cordgrass	Rau and Fassliotis, 1965
<b>61</b>	<i>Mesoanguina plantaginis</i>	Bracted plantain	Vargas and Sasser, 1976
<b>62</b>	<i>Mesocriconema curvatum</i>	Bermuda grass, creeping bentgrass, creeping grasses, soybean	Zeng et al., 2012b
<b>63</b>	<i>M. ornatum</i>	Corn, blueberry, grasses, ornamentals, peanut	Ratanaworabhan and Smart, 1970; Barker et al., 1982; Schmitt and Barker, 1988; Powers et al., 2004; Jagdale et al., 2013
<b>64</b>	<i>M. sphaerocephalum</i>	Corn, peanut	López et al., 2012; Zeng et al., 2012b
<b>65</b>	<i>M. xenoplax</i>	Bermuda grass, Chinese holly, creeping bentgrass, ornamentals, peach	Aycock et al., 1976; Zeng et al., 2012b; Powers et al., 2004
<b>66</b>	<i>Nanidorus minor</i>	American holly, azalea, bentgrass, bermuda grass, boxwood, camellia, centipedegrass, Chinese holly, corn, cotton, creeping bentgrass, holly, loblolly pine, longleaf pine, ornamentals, peach, peanut, slash pine, soybean, tall fescue, tobacco, white clover	Schmitt and Barker, 1988; Zeng et al., 2012b; Huang et al., 2018
<b>67</b>	<i>Ogma floridense</i>	Bermuda grass	Zeng et al., 2012b
<b>68</b>	<i>Paratrichodorus allius</i>	Creeping bentgrass	Zeng et al., 2012b
<b>69</b>	<i>P. porosus</i>	Camellia, corn, sorghum, soybean	Huang et al., 2018
<b>70</b>	<i>Paratylenchus goldeni</i>	Boxwood, centipedegrass	Raski, 1975; Zeng et al., 2012b
<b>71</b>	<i>Pratylenchus brachyurus</i>	Cotton, peach, peanut, soybean, tobacco	Schmitt and Barker, 1988
<b>72</b>	<i>P. coffeae</i>	Peach	Grand, 1985
<b>73</b>	<i>P. macrostylus</i>	Fraser fir, red spruce	Hartman and Eisenback, 1991

74	<i>P. penetrans</i>	Bermuda grass, boxwood, peach, potato, soybean, tobacco	Grand, 1985; Zeng et al., 2012b
75	<i>P. pratensis</i>	Cotton, ornamentals, potato, tobacco	Steiner, 1938; Grand, 1985
76	<i>P. scribneri</i>	Common bean, peach, soybean, tobacco	Grand, 1985
77	<i>P. vulnus</i>	Boxwood, Chinese holly, peach	Society of Nematologists, 1984; Grand, 1985
78	<i>P. zae</i>	Corn, peach, tall fescue	Grand, 1985; Schmitt and Barker, 1988
79	<i>Quinislucius capitatus</i>	Unknown	Society of Nematologists, 1984
80	<i>Rotylenchulus reniformis</i>	Cotton, sweet potato	Grand, 1985; Koenning et al., 2004
81	<i>Rotylenchus buxophilus</i>	Azalea, boxwood	Grand, 1985
82	<i>R. pumilus</i>	Trees	Ruehle, 1968
83	<i>Scutellonema brachyurus</i>	Bermuda grass, corn, cotton, tobacco	Zeng et al., 2012b
84	<i>Trichodorus elefjohnsoni</i>	Tulip poplar	Bernard, 1992
85	<i>T. obtusus</i>	Bermuda grass, St. Augustine grass, zoysiagrass	Ye et al., 2015; Huang et al., 2018
86	<i>Trophurus</i> spp.	Azalea	Grand, 1985
87	<i>Tylenchorhynchus claytoni</i>	American holly, azalea, bermuda grass, camellia, centipedegrass, Chinese holly, common bean, corn, cotton, creeping bentgrass, lima bean, loblolly pine, longleaf pine, ornamentals, peach, peanut, potato, slash pine, soybean, strawberry, tall fescue, tobacco, tomato, white clover	Steiner, 1937; Ruehle and Sasser, 1962; Aycock et al., 1976; Schmitt and Barker, 1988; Zeng et al., 2012b
88	<i>T. maximus</i>	Grasses, tomato	Grand, 1985
90	<i>Xenocriconemella macrodora</i>	Boxelder maple	López et al., 2012
91	<i>Xiphinema americanum</i> <i>sensu lato</i>	American holly, azalea, bermuda grass, boxwood, centipedegrass, Chinese holly, corn, cotton, loblolly pine, ornamentals, peach, peanut, slash pine, soybean, strawberry, sweet potato, tall fescue, tobacco, tomato	Ruehle and Sasser, 1962; Zeng et al., 2012b
92	<i>Xiphinema bakeri</i>	Bermuda grass	Zeng et al., 2012b
93	<i>Xiphinema chambersi</i>	Azalea, bermuda grass, centipedegrass, ornamentals, rose, soybean	Ye, 2012; Zeng et al., 2012b
94	<i>Xiphinema krugi</i>	Bermuda grass, boxwood, cotton, ornamentals, soybean, tomato	Ye and Robbins, 2010

## References

- Aycock, R., Barker, K.R. and Benson, D.M. (1976). Susceptibility of Japanese holly to *Criconemoides xenoplax*, *Tylenchorhynchus claytoni*, and certain other plant-parasitic nematodes. *Journal of Nematology* 8, 26-31.
- Baldwin, J.G. and Sasser, J.N. (1979). *Meloidogyne megatyla* n. sp. a root-knot nematode from loblolly pine. *Journal Nematology* 11(1), 47-56.
- Barker, K.R. and Sasser, J.N. (1959). Biology and control of the stem nematode, *Ditylenchus dipsaci*. *Phytopathology* 49 (10), 664-670.
- Barker, K.R., Schmitt, D.P. and Campos, V.P. (1982). Response of peanut, corn, tobacco, and soybean to *Criconemella ornata*. *Journal of Nematology* 14, 576-581.
- Bernard, E.C. (1992). *Trichodorus elefjohnsoni* n. sp. (Nemata: Trichodoridae) from undisturbed Appalachian forest. *Journal of Nematology* 24, 78-83.
- Brzeski, M.W. (1974). Taxonomy of Hemicycliophorinae (Nematoda, Tylenchida). *Zeszyty Problemowe Postępów Nauk Rolniczych* 154, 237-330.
- Eisenback, J.D. (1982). Description of the blueberry root-knot nematode. *Meloidogyne carolinensis* n. sp. *Journal of Nematology* 14, 303-317.
- Golden, A.M., Rau, G.J. and Cobb, G.S. (1962). *Heterodera cyperi* (Heteroderidae), a new species of cyst-forming nematode. *Proceedings of the Helminthological Society of Washington* 29, 168-173.
- Grand, L.F. (1985). North Carolina Plant Disease Index. Issue 240 (Revised) of Technical Bulletin. North Carolina Agricultural Research Service. North Carolina State University. Raleigh, North Carolina. 157pp.
- Hartman, K.M. and Eisenback, J.D. (1991). Amended description of *Pratylenchus macrostylus* Wu, 1971 with SEM observations. *Journal of Nematology* 23, 104-109.
- Hirschmann, H. and Sasser, J.N. (1955). On the occurrence of an intersexual form in *Ditylenchus triformis* n. sp. (Nematoda: Tylenchida). *Proceedings of the Helminthological Society of Washington* 22, 115-123.
- Holguin, C.M., Baeza, J.A., Mueller, J.D. and Agudelo, P. (2015a). High genetic diversity and geographic subdivision of three lance nematode species (*Hoplolaimus* spp.) in the United States. *Ecology and Evolution* 5 (14), 2929-2944. <http://doi.org/10.1002/ece3.1568>
- Holguin, C.M., Mueller, J., Khalilian, A. and Agudelo, P. (2015b). Population dynamics and spatial distribution of Columbia lance nematode in cotton. *Applied Soil Ecology* 95, 107-114.
- Holguin, C.M., Ma, X., Mueller, J.D. and Agudelo, P. (2016). Distribution of *Hoplolaimus* species in soybean fields in South Carolina and North Carolina. *Plant Disease* 100 (1), 149-153.
- Huang, D., Yan, G., Gudmestad, N., Whitworth, J.L., Frost, K., Brown, C., Ye, W., Agudelo, P. and Crow, W.T. (2018). Molecular characterization and identification of stubby-root nematode species from multiple states in the United States. *Plant Disease* (In press). <https://doi.org/10.1094/PDIS-10-17-1668-RE>
- Jagdale, G., Holladay, T., Brannen, P., Cline, W., Agudelo, P., Nycepir, A. & Noe, J. (2013). Incidence and pathogenicity of plant-parasitic nematodes associated with blueberry (*Vaccinium* spp.) replant disease in Georgia and North Carolina. *Journal of Nematology* 45, 92-98.
- Koenning, S.R. (2004). Resistance of soybean cultivars to field populations of *Heterodera glycines* in North Carolina. *Plant Disease* 88, 942 - 950.
- Koenning, S.R., Kirkpatrick, T.L., Starr, J.L., Wrather, J.A., Walker, N.R. and Mueller, J.D. (2004). Plant-parasitic nematodes attacking cotton in the United States - Old and emerging production challenges. *Plant Disease* 88, 100-113.
- Kohl, L.M., Warfield, C.L., Benson, D.M. (2010). Population dynamics and dispersal of *Aphelenchoides fragaria* in nursery-grown lantana. *Journal of Nematology* 42, 332-341.
- Komm, D.A., Reilly, J.J. and Elliott, A.P. (1983). Epidemiology of a tobacco cyst nematode (*Globodera tabacum*) in Virginia. *Plant Disease* 67, 1249-1251.
- Lewis, S.A. and Golden, A.M. (1981). Description and SEM observations of *Dolichodorus marylandicus* n. sp. with a key to species of *Dolichodorus*. *Journal of Nematology* 13, 128-135.
- Loof, P.A.A. (1991). The family Pratylenchidae Thorne 1949. In Nickle, W.R., ed. *Manual of Agricultural Nematology*. Marcel Dekker, Inc., NY. pp. 363-422.
- López, M.A.C., Robbins, R.T. and Szalanski, A.L. (2012). Taxonomic and molecular identification of *Bakernema*, *Criconema*, *Hemicriconemoides*, *Ogma* and *Xenocriconemella* species (Nematoda: Criconematidae). *Journal*

of Nematology 44, 427-446.

López, M.A.C., Robbins, R.T. and Szalanski, A.L. (2013). Taxonomic and molecular identification of *Hemicaloosia*, *Hemicyclophora*, *Gracilaculus* and *Paratylenchus* species (Nematoda: Criconematidae). Journal of Nematology 45, 145-171.

McCuston, J.L., Hudson, L.C., Subbotin, S.A., Davis, E.L. and Warfield, C.Y. (2007). Conventional and PCR detection of *Aphelenchoides fragariae* in diverse ornamental host plant species. Journal of Nematology 39 (4), 343-355.

Melton, T.A., Phillips, J.A., Imbriani, J.L. and Barker, K.R. (1991). First report of *Globodera tabacum solanacearum* on flue-cured tobacco outside Virginia. Plant Disease 75, 1074.

NCDA&CS Agricultural Statistics. (2017). 2017 Annual Statistics Book. <http://www.ncagr.gov/stats/AgStat/NCAgStatBook.pdf>

Nicol, J.M., Turner, S.J., Coyne, D.L., den Nijs, L., Hockland, S. and Maafi, Z.T. (2011). Current nematode threats to world agriculture. In: Genomics and Molecular Genetics of Plant—Nematode Interactions (Jones, J.T., Gheysen, G. and Fenoll, C., eds). Heidelberg: Springer. pp. 21-44.

Powers, T.O., Bernard, E.C., Harris, T., Higgins, R., Olson, M., Lodema, M., Mullin, P., Sutton, L. and Powers, K.S. (2014). COI haplotype groups in *Mesocriconema* (Nematoda: Criconematidae) and their morphospecies associations. Zootaxa 3827, 101-146.

Raski, D.J. (1975). Revision of the genus *Paratylenchus* Micoletzky, 1922 and descriptions of new species. Part 1 of 3 parts. Journal of Nematology 7, 15-34.

Ratanaworabhan, S. and Smart, Jr. G.C. (1970). The ring nematode, *Criconemoides ornatus*, on peach and centipede grass. Journal of Nematology 2 (3), 204-208.

Rau, G.J. and Fassuliotis, G. (1965). *Hypsoperine spartinae* n. sp., a gall-forming nematode on the roots of smooth cordgrass. Proceedings of the Helminthological Society of Washington 32, 159-162.

Ruehle, J.L. (1968). Distribution of plant-parasitic nematodes associated with forest trees of the world. U.S. Forestry Services. Southeast Forest Experimental Station, Ashville, North Carolina.

Ruehle, J.L. and Sasser, J.N. (1962). The role of plant-parasitic nematodes in stunting of pines in southern plantations. Phytopathology 52 (1), 56-68.

Schmitt, D.P and Barker, K.R. (1988). Incidence of plant-parasitic nematodes in the coastal plain of North Carolina. Plant Disease 72, 107-110.

Sher, S.A. (1963) Revision of the Hoplolaiminae (Nematoda). II. *Hoplolaimus* Daday, 1905 and *Aorolaimus* n. gen. Nematologica 9, 267-295.

Steiner, G. (1937) Opuscula miscellana nematologica. V. Proceedings of the Helminthological Society of Washington 4, 33-38.

Strider, D.L. (1979) Control of *Aphelenchoides ritzemabosi* in African violet. Plant Disease Reporter 63, 378-382.

Thorne, G. (1955). Fifteen new species of the genus *Hemicyclophora* with an emended description of *H. typica* de Man (Tylenchida: Criconematidae). Proceedings of the Helminthological Society of Washington 22, 1-16.

Triantaphyllou, A.C. and Hirschmann, H. (1973). Environmentally controlled sex expression in *Meloidodera floridensis*. Journal of Nematology 5, 181-185.

Van den Berg, E., Tiedt, L.R., Liébanas, G., Chitambar, J.J., Stanley, J.D., Inserra, R.N., Castillo, P. and Subbotin, S.A. 2018. Morphological and molecular characterisation of two new *Hemicyclophora* species (Tylenchida: Hemicyclophoridae) with a revision of the taxonomic status of some known species and a phylogeny of the genus. Nematology 20 (4), 319-354.

Vargas, O.F. and Sasser, J.N. (1976). Biology of *Anguina plantaginis* parasitic on *Plantago aristata*. Journal of Nematology 8 (1), 64-68.

Winstead, N. N. and Sasser, J. N. (1956). Reaction of cucumber varieties to five root-knot nematodes (Meloidogyne spp.). Plant Disease Reporter 40, 272-275.

Ye, W. and Giblin-Davis, R.M. (2013). Molecular characterization and development of real-time PCR assay for pine-wood nematode *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae). PLoS ONE 8 (11). November 24, 2015. DOI: 10.1371/journal.pone.0143556

Ye, W. and Robbins, R.T. (2000). Morphology of four species of *Hemicriconemoides* (Nematoda: Criconematidae) in the USA with the synonym of *H. annulatus*. International Journal of Nematology 10, 101-111.

Ye, W. and Robbins, R.T. (2010). Morphology and taxonomy of *Xiphinema* (Nematoda: Longidoridae) occurring in Arkansas, USA. Acta Agriculturae Universitatis Jiangxiensis 32 (5), 928-945.

Ye, W. (2012). Development of PrimeTime-Real-Time PCR for species identification of soybean cyst nematode (*Heterodera glycines* Ichinohe, 1952) in North Carolina. Journal of Nematology 44 (3), 284–290.

Ye, W. (2017). Soybean cyst nematode (*Heterodera glycines*) distribution in North Carolina, U.S.A. Plant Health Progress 18, 230-232. doi:10.1094/PHP-08-17-0050-BR

Ye, W., Koenning, S.R., Zhuo, K. and Liao, J. (2013). First report of *Meloidogyne enterolobii* on cotton and soybean in North Carolina, USA. Plant Disease 97 (9), 1262.

Ye, W., Zeng, Y., Tredway, L., Martin, S. and Martin, M. (2012). Plant parasitic nematodes in Carolina turfgrass. Carolina Green. March/April: 26 - 28. [http://spectrumcreativegraphics.com/carolinagreen\\_marapr12/](http://spectrumcreativegraphics.com/carolinagreen_marapr12/)

Ye, W., Zeng, Y. and Kerns, J. (2015a). First report of *Trichodorus obtusus* on turfgrass in North Carolina, USA. Plant Disease 99 (2), 291. <http://dx.doi.org/10.1094/PDIS-08-14-0830-PDN>

Ye, W., Zeng, Y. and Kerns, J. (2015b). Molecular characterization and diagnosis of root-knot nematodes (*Meloidogyne* spp.) from turfgrasses in North Carolina, USA. PLoS ONE 10 (11), e0143556. doi: 10.1371/journal.pone.0143556

Zeng, Y., Ye, W., Kerns, J., Martin, S., Tredway, L. and Martin, M. (2015). Molecular characterization and phylogenetic relationships of plant parasitic nematodes associated with turfgrasses in North Carolina and South Carolina, USA. Plant Disease 99 (7), 982-993.

Zeng, Y., Ye, W., Tredway, L., Martin, S. and Martin, M. (2012a). Description of *Hemicaloosia graminis* n.sp. (Nematoda: Caloosidae) associated with turfgrasses in North and South Carolina, USA. Journal of Nematology 44 (2), 134-141.

Zeng, Y., Ye, W., Tredway, L., Martin, S. and Martin, M. (2012b). Taxonomy and morphology of plant parasitic nematodes associated with turf grasses in North and South Carolinas, USA. Zootaxa 3452, 1-46.