Fraser Fir Seed Stratification

Protocol for germinating *Abies fraseri* seed

1. Introduction

Fraser fir (*Abies fraseri*) seeds exhibit a profound, deep physiological embryo dormancy common to many high-elevation conifers. In nature, this mechanism prevents autumn-shed seeds from germinating prematurely during transient warm spells before winter, safeguarding the next generation from fatal frost damage. To break this dormancy, the seed requires an extended period of low-temperature, highly oxygenated, moist conditions. We replicate this process artificially via the cold stratification process explained below.

2. The Value of Stratification

For Christmas tree growers and commercial nurseries, the direct value of proper stratification lies in maximizing efficiency, physical bench utilization, and crop consistency. Unstratified or poorly stratified Fraser fir seeds demonstrate erratic, slow, and highly protracted germination windows that can stretch across 60 to 90 days. Stratification addresses these issues with several key advantages:

KEY ADVANTAGES:

- **Synchronized Emergence:** Stratification condenses the germination window from weeks to days. This yields a highly uniform seedling canopy, ensuring that all individual units undergo subsequent culturing steps (fertilization, pest management, shading changes) at identical developmental milestones.
- **Improved Germination Capacity:** Stratification maximizes the biological potential of the lot, raising final germination rates by up to 40% compared to untreated controls.
- **Broadened Thermal Window:** Stratified seeds exhibit resilience to a wider range of seedbed temperatures, minimizing seedling losses during unpredictable spring microclimatic swings.

3. Step-by-Step Stratification Protocol

01



HYDRATION

Submerge the dry seed lot entirely in a clean vessel filled with room-temperature, distilled, or highly filtered water. The volume of water should be at least four times the volume of the seed lot to ensure adequate rehydration.

Allow the seeds to soak for 24 hours.

During this timeframe, viable seeds absorb water and sink. Blighted, insect-damaged, or empty seed coats will remain floating on the surface. Skim off and discard these floaters to further improve germination rate

02



STRATIFICATION MEDIUM MIX

Thoroughly drain the imbibed seeds. Prepare a clean stratification medium such as below:

- 1:1 mixture of horticultural-grade peat moss and coarse perlite (or vermiculite). Moisten the medium with water and mix thoroughly.
- Dampen a paper towel, squeeze out the excess moisture
- No stratification medium

Thoroughly fold the drained seeds uniformly into the damp medium.

03

Transfer the seed-and-medium mixture (or drained seed if not using medium) into heavy-duty poly zip-lock bags. Close



COLD CHILL

the bags leaving a small, finger-sized opening to allow for a small amount of airflow.

Place the sealed bags inside a dark refrigeration unit set between 1°C – 4°C (34°F – 39°F).

Keep them chilled for 6 to 8 weeks (42 to 60 days). Check the seed weekly for moisture, adding water if the seed dries out. This extended cold period is essential for breaking their dormancy.

04



SOWING

After a minimum of 42 days of chilling the Fraser fir seeds are ready for sowing. Stratified Fraser fir seed should germinate within 2-3 weeks, post-sowing.

4. Advanced Techniques

The Stratification-Redry Method

A persistent challenge with conventional stratification is that once the 60-day cycle concludes, the seeds are metabolically active and must be planted immediately. If unfavorable field or greenhouse conditions delay sowing, the seed lot may spoil or undergo premature "in-bag" germination, rendering them useless for mechanical seeders.

To mitigate this risk, some seed centers utilize the “Stratification-Redry” Method. Under this advanced protocol, the seeds are initially subjected to standard cold moist stratification for a slightly condensed duration (typically 4 weeks). Upon removal, the surface moisture is gently evaporated using clean fans, and the seeds are intentionally dried back to an intermediate, stable target moisture content of 25% to 30%.

Once dried back, the metabolic acceleration halts, locking the seed into a state of "suspended preparation." These seeds can then be held safely in cold dry storage (-1°C to 2°C) for several months. When finally sown into warm seedbeds, they retain their broken dormancy status and emerge rapidly and uniformly.

5. Additional Literature & Technical References

Adkins, C. R., Hinesley, L. E., & Blazich, F. A. (1984). Role of stratification, temperature, and light in germination of Fraser fir seed. Canadian Journal of Forest Research, 14(1), 88-93.

Bonner, F. T., & Karrfalt, R. P. (Eds.). (2008). The Woody Plant Seed Manual. Agriculture Handbook No. 727. Washington, DC: U.S. Department of Agriculture, Forest Service. (Section on Abies Mill. genus).

Leadem, C. L. (1986). Stratification-redry treatments to improve the germination of conifer seeds. Tree Planters' Notes, 37(3), 33-37.

Speers, C. F. (1962). Fraser fir seed collection, stratification, and germination. Tree Planters' Notes, 53, 19-21.